

On the probability for subhalo detection through quasar image-splitting

T. Riehm¹, E. Zackrisson^{1,2,3},
 K. Wiik² and O. Möller⁴

¹Stockholm Observatory, AlbaNova University Center, 106 91 Stockholm, Sweden

²Tuorla Observatory, University of Turku, Väisäläntie 20, FI-21500 Piikkiö, Finland

³Department of Astronomy and Space Physics, Box 515, 751 20 Uppsala, Sweden

⁴Max Planck Institut für Astrophysik, D-85741 Garching, Germany

Abstract. Dark galaxies may in principle be detectable through strong-lensing image splitting of quasars on small angular scales (milliarcseconds or below). Here, we estimate the overall probabilities for such detections under the assumption that the quasars can be treated as point sources. Due to the very low probabilities derived, we conclude that it is currently not feasible to use this strategy to put the CDM predictions for the dark galaxy population to the test.

Keywords. dark matter - cosmology: observations - galaxies: halos - quasars: general

1. CDM subhalos

In the CDM scenario (as well as in several slight modifications thereof), dark matter halos are assembled hierarchically from smaller subunits. A long-standing problem with this picture is that the number of subhalos predicted by CDM simulations at any given time is orders of magnitudes higher than the known number of satellite galaxies in the vicinity of the Milky Way. There are several possible ways out of this dilemma: either the CDM scenario is incorrect, the simulation predictions are wrong, or the majority of these subhalos must somehow have evaded detection. The latter alternative is quite viable, provided that the baryonic content of these subhalos has either been lost or prevented to form stars. If such “dark galaxies” do indeed exist, gravitational lensing may offer one of the most promising ways to detect them.

2. Gravitational millilensing by subhalos

One tell-tale signature of dark matter subhalos in the 10^6 – 10^{10} M_\odot range would be gravitational millilensing, i.e. image splitting with a characteristic separation of milliarcseconds (e.g. Yonehara et al. 2003). Based on a null detection of millilensing in a sample of 300 quasars observed with the VLBI, Wilkinson et al. (2001) demonstrated that the vast majority of quasars do not show any signs of millilensing, and were able to impose upper limits of $\Omega < 0.01$ on the cosmological density of dark matter objects in this mass range. Unfortunately, this constraint is still insufficient to rule out the subhalos predicted by CDM, since their optical depth is expected to be at least one order of magnitude lower. To put the CDM subhalo predictions to the test, Yonehara et al. (2003) instead suggested that one should target quasars which are already known to be gravitationally lensed on arcsecond scales, as one can then be sure that there is a massive halo well-aligned with the line of sight, which substantially increases the probability for subhalo millilensing. Indeed, the magnification associated with millilensing has long been suspected to be

the cause of the flux ratio anomalies seen in such systems (e.g. Mao & Schneider 1998; Kochanek & Dalal 2004). Subhalo millilensing has also been advocated as an explanation for strange bending angles of radio jets (Metcalf 2002) and image positions which smooth halo models seem unable to account for (Biggs et al. 2004).

3. Prospects for direct detection of dark matter subhalos through gravitational lensing

Under the assumption that the lenses do not overlap along the sightline, the optical depth τ represents the fraction of the sky that is covered by regions in which a point source will be lensed. In the limit of small τ , the optical depth can therefore directly be used as an estimate of the lensing probability. Assuming dark matter halos to be Singular Isothermal Spheres (SIS) and using the subhalo mass function and number density profile given in Gao et al. (2004), we derived estimates for the optical depth τ as a function of impact parameter to the center of the host halo. Even when allowing for a variety of host halo masses and redshift combinations, τ does not exceed a value of 10^{-3} . This is contrary to previous estimates (e.g. Yonehara et al. 2003) indicating optical depths several magnitudes higher. In order to be able to use this method to put the CDM subhalo predictions to a test, a survey of more than 10^3 objects would be required, in contrast to the $\sim 10^2$ multiply-imaged quasars presently known. Moreover, since it would be highly time consuming to observe on the order of 10^3 objects at milliarcsecond resolution using existing facilities, a search for CDM subhalos is currently not feasible. Furthermore, using SIS cross sections results only in an upper limit for the optical depth. Considering more realistic, less concentrated halo profiles, e.g. a Navarro-Frenk-White- (Navarro et al. 1996) or Moore-profile (Moore et al. 1999), would reduce the overall probabilities for detecting subhalos through this technique even more (Zackrisson et al. 2007). These results are valid for point sources. Whether extended sources (as in the case of VLBI observations of quasars) can boost the probability for subhalo millilensing is something that we are currently looking into (Riehm et al., in preparation).

Acknowledgements

EZ acknowledges research grants from the Swedish Research Council, the Academy of Finland and the Swedish Academy of Sciences.

References

- Biggs, A.D., Browne, I.W.A., Jackson, N.J., York, T., Norbury, M.A., McKean, J.P. & Phillips, P.M. 2004, *MNRAS* 350, 949
- Gao, L., White, S.D.M., Jenkins, A., Stoehr, F. & Springel, V. 2004, *MNRAS* 355, 819
- Kochanek, C.S. & Dalal, N. 2004, *ApJ* 610, 69
- Mao, S. & Schneider, P. 1998, *MNRAS* 295, 587
- Metcalf, R.B. 2004, *ApJ* 580, 696
- Moore, B., Quinn, T., Governato, F., Stadel, J. & Lake, G. 1999, *MNRAS* 310, 1147
- Navarro, J.F., Frenk, C.S. & White, S.D.M. 1996, *ApJ* 462, 563
- Wilkinson, P.N., et al. 2001, *Phys. Rev. Lett.* 86, 584
- Yonehara, A., Umemura, M. & Susa, H. 2003, *PASJ* 55, 1059
- Zackrisson, E., Riehm, T., Möller, O., Wiik, K. & Nurmi, P. 2007, submitted to *ApJ*